



# Dimension<sup>®</sup> 0.25% + Fertilizer

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes, and sod farms.

Covers up to 41,660 sq ft (100'x417')

In the state of New York, this product may be applied only by commercial applicators at no more than 200 lb (0.5 lb active ingredient) per acre (4.6 lb per 1,000 sq ft) per year. Use of this product in Nassau and Suffolk Counties of New York is prohibited.

### ACTIVE INGREDIENT:

Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester .....0.25%

OTHER INGREDIENTS.....99.75%

Total:.....100.0%

This product contains 0.125 pound of the active ingredient dithiopyr per 50 pound bag.

# 18-0-3

## GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N) .....18.0%  
18.0% Urea Nitrogen\*

SOLUBLE POTASH (K<sub>2</sub>O) .....3.0%

SULFUR (S) .....1.4%  
1.4% Free Sulfur (S)

DERIVED FROM: Urea, Polymer-Coated Sulfur-Coated Urea, Muriate of Potash.

CHLORINE (Cl) Max. ....2.3%

\* 4.0% Slowly Available Nitrogen from Polymer-Coated Sulfur-Coated Urea.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### Precautionary Statements

Hazards to Humans and Domestic Animals  
Causes Eye Irritation • Harmful if Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

### Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove clothing/PPE if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### FIRST AID

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, the give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

**Environmental Hazards:** This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Refer to back and front of bag for Directions for Use including Storage and Disposal.

**Notice:** Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

**Agricultural Chemical: Do not ship or store with food, feed, drugs or clothing.**

### Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Agricultural Use Requirements:** Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

**Non-Agricultural Use Requirements:** The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

- Keep unprotected persons out of treated area during application and until dust has settled.

### Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area.

**Pesticide Disposal:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Use Directions for Turf

TCS GrowStar Dimension<sup>®</sup> 0.25% + Fertilizer provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes, and sod farms. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

### Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury.

• Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/gfbbmp07.pdf>.

• Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at [http://www.dep.state.fl.us/central/Home/Meetings/Training/FLGreen/BMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/central/Home/Meetings/Training/FLGreen/BMP_Book_final.pdf).

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses	
Bentgrass, creeping *	<i>Agrostis palustris</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, fine **	<i>Festuca rubra</i>
Fescue, tall	<i>Festuca arundinacea</i>
Ryegrass, perennial	<i>Lolium perenne</i>
Warm-Season Grasses	
Bahiagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss ***	<i>Buchloe dactyloides</i>
Carpetgrass	<i>Axonopus affinis</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Colonial bentgrass (*Agrostis tenuis*).

\*\* Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\* Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

**Note:** it is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

### Precautions for Reseeding, Overseeding, or Sprigging

• Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 4.6 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.

• When reseeded or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• For best overseeding results, use mechanical or power seeding equipment (slit seeders) designed to provide good contact between soil and seed.

### Precautions for Control of Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

### Other Use Precautions

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from area treated with this product.

# Net Weight: 50 LBS (22.67 KG) Part #: 901528



# Dimension® 0.25% + Fertilizer

## Application Directions for Turf

### Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
  - The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
  - Use of split (half-rate) applications spaced 5 - 10 weeks apart may provide improved weed control.
- Maximum Use Rates (Applies to all Turfgrass Uses)
- Do not apply more than 200 lb (0.5 lb active ingredient) per acre (4.6 lb per 1000 sq ft) per application or more than 600 lb (1.5 lb active ingredient) per acre (13.8 lb per 1000 sq ft) per year if using split or sequential applications.
  - In the state of New York, this product may be applied only by commercial applicators at no more than 200 lb (0.5 lb active ingredient) per acre (4.6 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

### Control of Crabgrass

#### Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides "early postemergence" control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2 - 4 week longer period of time (window of application) during which crabgrass control can be achieved.

#### Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described in the following use programs.

**Spring Application:** For applications made in the spring or early summer, use an appropriate application rate corresponding to one of the three control programs listed in Table 1. Specific rate recommendations are based on the user's location (Region), the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

**Use Program 1** for preemergence control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3 - 5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

**Use Program 2** for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., residential lawns).

**Use Program 3** for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively low (e.g., golf fairways).

Table 1 Recommended Use Rates

Region	Application Rates	Program 1	Program 2	Program 3
All states, except NY, and parts of states not included in transition, south, coastal south or west.	lb/acre	52	100	152
	lb/1000 sq ft	1.2	2.3	3.5
	lb ai/acre	0.13	0.25	0.38
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI	lb/acre	100	152	200
	lb/1000 sq ft	2.3	3.5	4.6
	lb ai/acre	0.25	0.38	0.5
South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	lb/acre	152	100 + 100	124 + 124
	lb/1000 sq ft	3.5	2.3 + 2.3	2.8 + 2.8
	lb ai/acre	0.38	0.25 + 0.25	0.31 + 0.31
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	lb/acre	100 + 100	124 + 124	152 + 152
	lb/1000 sq ft	2.3 + 2.3	2.8 + 2.8	3.5 + 3.5
	lb ai/acre	0.25 + 0.25	0.31 + 0.31	0.38 + 0.38
West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	lb/acre	76 + 100	100 + 152	100 + 100
	lb/1000 sq ft	1.7 + 2.3	2.3 + 3.5	2.3 + 2.3
	lb ai/acre	0.19 + 0.25	0.25 + 0.38	0.25 + 0.25

\* Split applications may be made 5 - 10 weeks apart.

**Use of Split or Sequential Applications:** Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with the rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above and separate rate restrictions for the state of New York.

**Fall Application:** This product can also be applied in the late summer through early fall (late August through November) at use rates listed in Table 1 for "Program 3" to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

## Weeds Controlled or Suppressed

**Spring Application:** When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds at application rates given in the Table 1 above:

Grasses		Broadleaf Weeds	
barley	<i>Hordeum</i> spp.	bittercress •	<i>Cardamine</i> spp.
barnyardgrass	<i>Echinochloa crus-galli</i>	carpetweed	<i>Mollugo verticillata</i>
bluegrass, annual	<i>Poa annua</i>	chickweed •	<i>Stellaria</i> spp.
brome	<i>Bromus</i> spp.	geranium, Carolina •	<i>Geranium carolinianum</i>
crabgrass, large	<i>Digitaria sanguinalis</i>	henbit	<i>Lamium</i> spp.
crabgrass, smooth	<i>Digitaria ischaemum</i>	knotted, prostrate	<i>Polygonum aviculare</i>
crabgrass, Southern	<i>Digitaria ciliaris</i>	lespedeza, common •	<i>Lespedeza striata</i>
crowfootgrass •	<i>Dactyloctenium aegyptium</i>	marestail	<i>Conyza canadensis</i>
dallisgrass (seedling)	<i>Paspalum dilatatum</i>	medic, black	<i>Medicago lupulina</i>
goosegrass	<i>Eleusine indica</i>	mustard	<i>Brassica</i> spp.
foxtail, green	<i>Setaria verdi</i>	oxalis, buttercup	<i>Oxalis pes-caprae</i>
foxtail, yellow	<i>Setaria faberi</i>	pineappleweed •	<i>Matricaria matricarioides</i>
kikuyugrass •	<i>Pennisetum clandestinum</i>	pigweed, redroot	<i>Amaranthus retroflexus</i>
oats, wild	<i>Avena fatua</i>	parsley-piert •	<i>Alchemilla arvensis</i>
ryegrass (annual & perennial)	<i>Lolium</i> spp.	purslane, common	<i>Portulaca oleracea</i>
sandbur	<i>Cenchrus</i> spp.	rocket, London	<i>Sisymbrium irio</i>
smutgrass	<i>Sporobolus indicus</i>	shepherdspurse	<i>Capsella bursa-pastoris</i>
		speedwell, corn •	<i>Veronica arvensis</i>
		spurge, garden	<i>Euphorbia hirta</i>
		spurge, prostrate	<i>Euphorbia humistrata</i>
		spurge, spotted	<i>Euphorbia maculata</i>
		woodsorrel, creeping	<i>Oxalis corniculata</i>
		woodsorrel, yellow	<i>Oxalis stricta</i>

\* Suppression only

**Fall Application:** When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabgrass in the following spring.

bluegrass, annual	<i>Poa annua</i>	geranium, Carolina	<i>Geranium carolinianum</i>
bittercress	<i>Cardamine</i> spp.	parsley-piert	<i>Alchemilla arvensis</i>
chickweed	<i>Stellaria</i> spp.	pineappleweed	<i>Matricaria matricarioides</i>
henbit	<i>Lamium</i> spp.	shepherdspurse	<i>Capsella bursa-pastoris</i>

### Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

### Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

### Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Dimension® is a Trademark of DOW AgroSciences LLC  
TCS GrowStar® is a Registered Trademark of Turf Care Supply Corp.

**Net Weight: 50 LBS (22.67 KG)**

EPA REG. NO. 62719-492-82757  
EPA EST. NO. 82757-OH-001 (MF)  
EPA EST. NO. 82757-MA-002 (H)  
EPA EST. NO. 82757-FL-001 (S)  
(First letter of lot # indicates Manufacturing Site)  
**#901528**

F1612

Distributed by:  
**Turf Care Supply Corp.**  
50 Pearl Road, Suite 200  
Brunswick, OH 44212

**TurfCare™**  
supply corp.